

VACCINE GUIDELINES

DISEASE/ VACCINE	FOALS/ WEANLINGS	BROODMARES	ADULT HORSES OVER 1 YEAR OF AGE PREVIOUSLY VACCINATED AGAINST DISEASE INDICATED	ADULT HORSES OVER 1 YEAR OF AGE UNVACCINATED OR LACKING VACCINATION HISTORY
ENCEPHALOMYELITIS (EEE, WEE, VEE) *Consider 6-month revaccination interval for immunocompromised horses or horses residing in endemic areas.	From vaccinated mares: 1st dose at 4 - 6 months of age; 2nd dose 4 - 6 weeks after 1st dose; 3rd dose at 10 - 12 months of age, prior to the onset of the next vector season From non-vaccinated mares: 1st dose at 3 - 4 months of age; 2nd dose 4 weeks after 1st dose; 3rd dose, 8 week interval after 2nd dose	Previously vaccinated: Annual, 4 - 6 weeks pre-partum. Previously unvaccinated or having unknown vaccination history 2-dose series: 2nd dose 4 weeks after 1st dose. Revaccinate 4 - 6 weeks prepartum.	Annual – spring, prior to onset of vector season.	2-dose series: 2nd dose 4 - 6 weeks after 1st dose. Revaccinate prior to the onset of the next vector season.
TETANUS TOXOID *Additional booster should be administered at time of penetrating wound or surgery if last dose is not within 6 months.	From vaccinated mares: 1st dose at 4 - 6 months of age; 2nd dose 4 - 6 weeks after the 1st dose; 3rd dose at 10 - 12 months of age. From non-vaccinated mares: 1st dose at 3 - 4 months of age; 2nd dose 4 - 6 weeks after the 1st dose; 3rd dose 10 - 12 months of age.	Previously vaccinated: Annual, 4 - 6 weeks pre-partum. Previously unvaccinated or having unknown vaccination history 2-dose series: 2nd dose 4 weeks after 1st dose. Revaccinate 4 - 6 weeks prepartum.	Annual	2-dose series: 2nd dose 4 - 6 weeks after 1st dose. Annual revaccination
WEST NILE VIRUS *Consider 6-month revaccination interval for geriatric, immunocompromised and horses residing in endemic areas.	From vaccinated mares: 1st dose at 4 - 6 months of age; 2nd dose 4 - 6 weeks after 1st dose; 3rd dose at 10 - 12 months of age, prior to the onset of the next vector season From non-vaccinated mares: 3-dose series: 1st dose at 3 - 4 months of age; 2nd dose, 4 weeks after 1st dose; 3rd dose, 8 week interval after 2nd dose.	Previously vaccinated: Annual, 4 - 6 weeks pre-partum. Unvaccinated or lacking vaccination history: It is preferable to vaccinate naïve mares when open.	Annual – spring, prior to onset of vector season	2-dose series: 2nd dose 4 - 6 weeks after 1st dose. Revaccinate prior to the onset of the next vector season.
INFLUENZA	*3-dose series: 1st dose at 6 months of age; 2nd dose 3 - 4 weeks after 1st dose; 3rd dose at 10 - 12 months of age. <i>*Both for vaccinated and unvaccinated mares.</i>	Previously vaccinated: Annual, 4 - 6 weeks pre-partum. Previously unvaccinated or having unknown vaccination history 3-dose series: 2nd dose 4 - 6 weeks after 1st dose; 3rd dose 4 - 6 weeks pre-partum.	Horses with ongoing risk of exposure: Semi-annual Horses at low risk of exposure: Annual	3-dose series: 2nd dose 4 - 6 weeks after 1st dose; 3rd dose 3 - 6 months after 2nd dose. Revaccinate semi-annually to annually
RHINOPNEUMONITIS (EHV-1 and EHV-4)	3-dose series: 1st dose 4 - 6 months of age; 2nd dose 4 - 6 weeks after 1st dose; 3rd dose at 10 - 12 months of age; Revaccinate at 6-month intervals. <i>*For foals from both vaccinated and unvaccinated mares.</i>	3-dose series <i>with product labeled for protection against EHV abortion:</i> Give at 5, 7 and 9 months of gestation	Annual Consider 6-month revaccination interval for: 1) Horses less than 5 years of age; 2) Horses on breeding farms or in contact with pregnant mares; 3) Performance or show horses at high risk.	3-dose series: 2nd dose 4-6 weeks after 1st dose; 3rd dose 4-6 weeks after 2nd dose.
RABIES	From vaccinated mares: 1st dose at 6 months of age; 2nd dose 4 - 6 weeks after 1st dose. From non-vaccinated mares: 1 dose Annual revaccination	Annual, 4 - 6 weeks pre-partum OR Prior to breeding, due to the relatively long duration of immunity, this vaccine may be given post-foaling but prior to breeding.	Annual	Single dose Annual revaccination
STRANGLES *Optional - recommended at locations where have had previous outbreaks, however vaccination is not recommended during an outbreak situation.	1st dose at 4 - 6 months of age; 2nd dose 4 - 6 weeks after 1st dose; 3rd dose 4 - 6 weeks after 2nd dose.	Previously vaccinated: Semi-annual with one dose given 4 - 6 weeks pre-partum. Previously unvaccinated or having unknown vaccination history: 3-dose series: 2nd dose 2 - 4 weeks after 1st dose; 3rd dose 4 - 6 weeks pre-partum.	Semi-annual to annual	2-3-dose series: 2nd dose 2 - 4 weeks after 1st dose; 3rd dose (where recommended by manufacturer) 2 - 4 weeks after 2nd dose. Revaccinate semi-annually

*Additional optional vaccinations (Botulism, Potomac Horse Fever and others) are available as needed.